

# VCC & VCF Collection 4 Reprocessing Status

MODLAND Collection 4
Reprocessing Review
17 December 2002
sohlberg@umd.edu


## Sequence of Presentation

- MOD44C 16-day interim 250m composite
- MOD44A Vegetative Cover Conversion
- MOD44B Vegetation Continuous Fields

# MOD44C: C4 Changes & Benefits

- •Incremental changes were made to Collection 3 thus differences between Collections 3 and 4 are minimal.
- •Improved utilization of Quality Flags:
  - Converted to MOD09 internal cloud flag;
  - •Refined use of aerosol flags;
  - •Instituted newly populated cirrus flag.
- •Refined the sorting routine:
  - •Improved format for ease of maintenance;
  - •Resolved one-off error in secondary layers table.
- •Collection 4 data continues efforts to select the best (cloud free, near nadir pixel) and will result in improved change detection.

#### Example from tile h12v10 (southern Amazon basin)



## MOD44C: C4 Known Issues

- •Dark features remain an issue that we would like to resolve:
  - •Features with spatial structure remain, particularly in arid areas, that have low reflectance relative to adjacent pixels.
  - •Possibly related to cloud shadow or over-correction.
  - •Will require ongoing investigation at the SCF.
  - •These low values are an issue because the values have characteristics consistent with forested areas.


# MOD44C: Anticipated Changes

- •Incremental refinements to the use of the cirrus flags are likely.
- •If we can find a resolution for the dark feature issue this will be submitted as soon as possible.
- •Update the QA flags that are retained to reflect current configuration.
- •Update production rule to accommodate case where the L2G compressed SDS is not filled (late season, high latitude case).

# MOD44A: C4 Changes & Benefits

- •With Collection 4, 250m data products will be available globally over the entire MODIS record.
- •As such, this will be the first opportunity to produce MOD44A results for large areas over an annual sample period.
- •SCF results with improved LUTs and composites are promising and we are quite eager to evaluate our change hotspot product at continental to global scales.
- •Primary C3 to C4 change is in the spectral LUTs, which to date we have only been able to derive globally at 500m resolution.

#### VCC Change Result: Mato Grosso Plateau, Brazil August 2001 to August 2002



## MOD44A: C4 Known Issues

- •Bug in output plane results in linear feature every n<sup>th</sup> line.
  - •Not certain if bug is in our code or the toolkit.
- •Need to alter production rule to add full Julian day to name of output file.
- •Update 16 to 32 day compositing method to reflect recent changes to MOD44C (cirrus and cloud flag).


## MOD44A: Anticipated Changes

- •Delivery of the revised LUTs.
- •Delivery of fixes for known issues from last graph.
- •Release of global, annual summary results.

## MOD44B: C4 Changes & Benefits

- •% Tree Cover to be distributed from LP-DAAC.
- •New layers being added:
  - •Leaf Characteristics
    - •% Broadleaf and % Needleleaf
  - Leaf Longevity
    - •% Evergreen and % Deciduous
  - •% Herbaceous
  - •%Bare
- •Much better ability to discern fragmented forest than AVHRR.


#### **MODIS Vegetation Continuous Fields – 2000 prototype**



Grass/shrubs/moss

Trees

Percent cover



## MOD44B: C4 Known Issues

•Collection 3 has a few areas of low signal due to persistent haze of clouds. Hopefully C4 surface reflectance improvements and the Aqua observations will help improve these troublesome areas.

# MOD44B: Anticipated Changes

- •Will produce annual products for 2000, 2001, and 2002.
- •Start dates will coincide with Spring Equinox.